Author Index Vol. 758

Açil, Y., Brinckmann, J., Behrens, P., Müller, P.K. and Bätge, B.
Semipreparative isolation of collagen types I, II, III and V by sodium dodecyl sulfate–polyacrylamide gel electrophoresis and electroelution 758(1997)313
Alexander, P.W., see Chen, Z. 758(1997)227
Alongi, D.M., see Susic, M. 758(1997)243
Antal, Jr., M.J., see Volgger, D. 758(1997)263
Arica, M.Y., see Denizli, A. 758(1997)217
Arrington, C.M., see Weldon, M.K. 758(1997)293
Bartle, K.D., see Johnson, B.R. 758(1997)65
Bartolozzi, F., Bertazza, G., Bassi, D. and Cristaferi, G.
Simultaneous determination of soluble sugars and organic acids as their trimethylsilyl derivatives in apricot fruits by gas–liquid chromatography 758(1997)99
Bassi, D., see Bartolozzi, F. 758(1997)99
Bätge, B., see Açil, Y. 758(1997)313
Behrens, P., see Açil, Y. 758(1997)313
Bensetiti, Z., see Yun, T. 758(1997)175
Berezkin, V.G. and Buzaev, V.V.
New thin-layer chromatography plate with a closed sorbent layer and details of its application 758(1997)125
Bergström, M., see Ohlson, S. 758(1997)199
Bertazza, G., see Bartolozzi, F. 758(1997)99
Blomberg, L.G., see Wan, H. 758(1997)303
Bonn, G.K., see Volgger, D. 758(1997)263
Borrego, F., see Montijano, H. 758(1997)163
Branch, S.K., Holzgrabe, U., Jefferies, T.M., Mallwitz, H. and Oxley, F.J.R.
Effect of β-cyclodextrin acetylation on the resolution of phenethylamines with capillary electrophoresis and nuclear magnetic resonance spectroscopy 758(1997)277
Brinckmann, J., see Açil, Y. 758(1997)313
Buzaev, V.V., see Berezkin, V.G. 758(1997)125
Campisi, B., see Damiani, P. 758(1997)109
Canales, I., see Montijano, H. 758(1997)163
Cantin-Esnault, D., see Oubrahim, H. 758(1997)145
Catterick, T., see Fairman, B. 758(1997)85
Channar, A.H., see Khuhawar, M.Y. 758(1997)159
Chase, H.A., see Nash, D.C. 758(1997)53
Chattopadhyay, A. and Hage, D.S.
Determination of the diol content of chromatographic supports by capillary electrophoresis 758(1997)255
Chen, Z. and Alexander, P.W.
Potentiometric detection of aliphatic amines by flow injection analysis and ion-interaction chromatography with a metallic copper electrode 758(1997)227
Cossignani, L., see Damiani, P. 758(1997)109
Cristaferi, G., see Bartolozzi, F. 758(1997)99
Damiani, P., Cossignani, L., Simonetti, M.S., Campisi, B., Favretto, L. and Favretto, L.G.
Stereospecific analysis of the triacylglycerol fraction and linear discriminant analysis in a climatic differentiation of Umbrian extra-virgin olive oils 758(1997)109
Denizli, A., Salih, B., Arica, M.Y., Kesenci, K., Hasirci, V. and Pişkin, E.
Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217
Fairman, B., Catterick, T., Wheals, B. and Polinina, E.
Reversed-phase ion-pair chromatography with inductively coupled plasma–mass spectrometry detection for the determination of organo-tin compounds in waters 758(1997)85
Favretto, L., see Damiani, P. 758(1997)109
Favretto, L.G., see Damiani, P. 758(1997)109
García, M.C., Torre, M., Laborda, F. and Marina, M.L.
Rapid separation of soybean globulins by reversed-phase high-performance liquid chromatography 758(1997)75
Gilliard, J.A. and Ritter, C.
Simulations of liquid chromatography–diode array detector data including instrumental artefacts for the evaluation of mixture analysis techniques 758(1997)1
Guiochon, G., see Yun, T. 758(1997)175
Hage, D.S., see Chattopadhyay, A. 758(1997)255
Hasirci, V., see Denizli, A. 758(1997)217
Herod, A.A., see Johnson, B.R. 758(1997)65
Holzgrabe, U., see Branch, S.K. 758(1997)277
Horie, H., Mukai, T. and Kohata, K.
Simultaneous determination of qualitatively important components in green tea infusions using capillary electrophoresis 758(1997)332
Jefferies, T.M., see Branch, S.K. 758(1997)277
Johnson, B.R., Bartle, K.D., Herod, A.A. and Kandiyoti, R.
N-Methyl-2-pyrrolidinone as a mobile phase in the size-exclusion chromatography of coal derivatives 758(1997)65
Kandiyoti, R., see Johnson, B.R. 758(1997)65
Kataoka, H., Matsumura, S., Koizumi, H. and Makita, M.
Rapid and simultaneous analysis of protein and non-protein amino acids as N(O,S)-isobutoxycarbonyl methyl ester derivatives by capillary gas chromatography 758(1997)167
Kesenci, K., see Denizli, A. 758(1997)217
Khuhawar, M.Y., Channar, A.H. and Lanjwani, S.N.
Indirect liquid chromatographic determination of malathion in formulations, based on the formation of palladium(II)–dimethyldithiosphate complex 758(1997)159
Kim, M., see Kiyohara, S. 758(1997)209
Kiyohara, S., Kim, M., Toida, Y., Saito, K., Sugita, K. and Sugo, T.
Selection of a precursor monomer for the introduction of affinity ligands onto a porous membrane by radiation-induced graft polymerization 758(1997)209
Kohata, K., see Horie, H. 758(1997)332
Koizumi, H., see Kataoka, H. 758(1997)167
Kozekov, I., see Palamareva, M. 758(1997)135
Laborda, F., see García, M.C. 758(1997)75
Lanjwani, S.N., see Khuhawar, M.Y. 758(1997)159
Lee, M.L., see Li, W. 758(1997)117
Li, W., Malik, A. and Lee, M.L.
Pressure drop effects in packed capillary column supercritical fluid chromatography 758(1997)117
Lorbeer, E., see Mayer, B.X. 758(1997)235
Lundblad, A., see Ohlson, S. 758(1997)199
Makita, M., see Kataoka, H. 758(1997)167
Malik, A., see Li, W. 758(1997)117
Mallwitz, H., see Branch, S.K. 758(1997)277
Marina, M.L., see García, M.C. 758(1997)75

Author Index Vol. 758

Açil, Y., Brinckmann, J., Behrens, P., Müller, P.K. and Bätge, B.
Semipreparative isolation of collagen types I, II, III and V by sodium dodecyl sulfate–polyacrylamide gel electrophoresis and electroelution 758(1997)313
Alexander, P.W., see Chen, Z. 758(1997)227
Alongi, D.M., see Susic, M. 758(1997)243
Antal, Jr., M.J., see Volgger, D. 758(1997)263
Arica, M.Y., see Denizli, A. 758(1997)217
Arrington, C.M., see Weldon, M.K. 758(1997)293
Bartle, K.D., see Johnson, B.R. 758(1997)65
Bartolozzi, F., Bertazza, G., Bassi, D. and Cristaferi, G.
Simultaneous determination of soluble sugars and organic acids as their trimethylsilyl derivatives in apricot fruits by gas–liquid chromatography 758(1997)99
Bassi, D., see Bartolozzi, F. 758(1997)99
Bätge, B., see Açil, Y. 758(1997)313
Behrens, P., see Açil, Y. 758(1997)313
Bensetiti, Z., see Yun, T. 758(1997)175
Berezkin, V.G. and Buzaev, V.V.
New thin-layer chromatography plate with a closed sorbent layer and details of its application 758(1997)125
Bergström, M., see Ohlson, S. 758(1997)199
Bertazza, G., see Bartolozzi, F. 758(1997)99
Blomberg, L.G., see Wan, H. 758(1997)303
Bonn, G.K., see Volgger, D. 758(1997)263
Borrego, F., see Montijano, H. 758(1997)163
Branch, S.K., Holzgrabe, U., Jefferies, T.M., Mallwitz, H. and Oxley, F.J.R.
Effect of β-cyclodextrin acetylation on the resolution of phenethylamines with capillary electrophoresis and nuclear magnetic resonance spectroscopy 758(1997)277
Brinckmann, J., see Açil, Y. 758(1997)313
Buzaev, V.V., see Berezkin, V.G. 758(1997)125
Campisi, B., see Damiani, P. 758(1997)109
Canales, I., see Montijano, H. 758(1997)163
Cantin-Esnault, D., see Oubrahim, H. 758(1997)145
Catterick, T., see Fairman, B. 758(1997)85
Channar, A.H., see Khuhawar, M.Y. 758(1997)159
Chase, H.A., see Nash, D.C. 758(1997)53
Chattopadhyay, A. and Hage, D.S.
Determination of the diol content of chromatographic supports by capillary electrophoresis 758(1997)255
Chen, Z. and Alexander, P.W.
Potentiometric detection of aliphatic amines by flow injection analysis and ion-interaction chromatography with a metallic copper electrode 758(1997)227
Cossignani, L., see Damiani, P. 758(1997)109
Cristaferi, G., see Bartolozzi, F. 758(1997)99
Damiani, P., Cossignani, L., Simonetti, M.S., Campisi, B., Favretto, L. and Favretto, L.G.
Stereospecific analysis of the triacylglycerol fraction and linear discriminant analysis in a climatic differentiation of Umbrian extra-virgin olive oils 758(1997)109
Denizli, A., Salih, B., Arica, M.Y., Kesenci, K., Hasirci, V. and Pişkin, E.
Cibacron Blue F3GA-incorporated macroporous poly(2-hydroxyethyl methacrylate) affinity membranes for heavy metal removal 758(1997)217
Fairman, B., Catterick, T., Wheals, B. and Polinina, E.
Reversed-phase ion-pair chromatography with inductively coupled plasma–mass spectrometry detection for the determination of organo-tin compounds in waters 758(1997)85
Favretto, L., see Damiani, P. 758(1997)109
Favretto, L.G., see Damiani, P. 758(1997)109
García, M.C., Torre, M., Laborda, F. and Marina, M.L.
Rapid separation of soybean globulins by reversed-phase high-performance liquid chromatography 758(1997)75
Gilliard, J.A. and Ritter, C.
Simulations of liquid chromatography–diode array detector data including instrumental artefacts for the evaluation of mixture analysis techniques 758(1997)1
Guiochon, G., see Yun, T. 758(1997)175
Hage, D.S., see Chattopadhyay, A. 758(1997)255
Hasirci, V., see Denizli, A. 758(1997)217
Herod, A.A., see Johnson, B.R. 758(1997)65
Holzgrabe, U., see Branch, S.K. 758(1997)277
Horie, H., Mukai, T. and Kohata, K.
Simultaneous determination of qualitatively important components in green tea infusions using capillary electrophoresis 758(1997)332
Jefferies, T.M., see Branch, S.K. 758(1997)277
Johnson, B.R., Bartle, K.D., Herod, A.A. and Kandiyoti, R.
N-Methyl-2-pyrrolidinone as a mobile phase in the size-exclusion chromatography of coal derivatives 758(1997)65
Kandiyoti, R., see Johnson, B.R. 758(1997)65
Kataoka, H., Matsumura, S., Koizumi, H. and Makita, M.
Rapid and simultaneous analysis of protein and non-protein amino acids as N(O,S)-isobutoxycarbonyl methyl ester derivatives by capillary gas chromatography 758(1997)167
Kesenci, K., see Denizli, A. 758(1997)217
Khuhawar, M.Y., Channar, A.H. and Lanjwani, S.N.
Indirect liquid chromatographic determination of malathion in formulations, based on the formation of palladium(II)–dimethyldithiosphate complex 758(1997)159
Kim, M., see Kiyohara, S. 758(1997)209
Kiyohara, S., Kim, M., Toida, Y., Saito, K., Sugita, K. and Sugo, T.
Selection of a precursor monomer for the introduction of affinity ligands onto a porous membrane by radiation-induced graft polymerization 758(1997)209
Kohata, K., see Horie, H. 758(1997)332
Koizumi, H., see Kataoka, H. 758(1997)167
Kozekov, I., see Palamareva, M. 758(1997)135
Laborda, F., see García, M.C. 758(1997)75
Lanjwani, S.N., see Khuhawar, M.Y. 758(1997)159
Lee, M.L., see Li, W. 758(1997)117
Li, W., Malik, A. and Lee, M.L.
Pressure drop effects in packed capillary column supercritical fluid chromatography 758(1997)117
Lorbeer, E., see Mayer, B.X. 758(1997)235
Lundblad, A., see Ohlson, S. 758(1997)199
Makita, M., see Kataoka, H. 758(1997)167
Malik, A., see Li, W. 758(1997)117
Mallwitz, H., see Branch, S.K. 758(1997)277
Marina, M.L., see García, M.C. 758(1997)75

Matsumura, S., see Kataoka, H. 758(1997)167
Mayer, B.X. and Lorbeer, E.
Triacylglycerol mixture for testing capillary columns for high-temperature gas chromatography 758(1997)235
McCreath, G.E., see Nash, D.C. 758(1997)53
Montijano, H., Borrego, F., Canales, I. and Tomás-Barberán, F.A.
Validated high-performance liquid chromatographic method for quantitation of neohesperidine dihydro-chalcone in foodstuffs 758(1997)163
Mukai, T., see Horie, H. 758(1997)332
Müller, P.K., see Açil, Y. 758(1997)313
Myers, P., see Zimina, T.M. 758(1997)191
Nash, D.C., McCreath, G.E. and Chase, H.A.
Modification of polystyrenic matrices for the purification of proteins. Effect of the adsorption of poly(vinyl alcohol) on the characteristics of poly(styrene–divinylbenzene) beads for use in affinity chromatography 758(1997)53
Ohlson, S., Bergström, M., Påhlsson, P. and Lundblad, A.
Use of monoclonal antibodies for weak affinity chromatography 758(1997)199
Okada, T.
Nonaqueous anion-exchange chromatography. I. Role of solvation in anion-exchange resin 758(1997)19
Okada, T.
Nonaqueous anion-exchange chromatography. II. Changing anion-exchange selectivity by resin surface complex formation of crown ethers 758(1997)29
Oubrahim, H., Richard, J.-M., Cantin-Esnault, D., Seigle-Murandi, F. and Trécourt, F.
Novel methods for identification and quantification of the mushroom nephrotoxin orellanine. Thin-layer chromatography and electrophoresis screening of mushrooms with electron spin resonance determination of the toxin 758(1997)145
Oxley, F.J.R., see Branch, S.K. 758(1997)277
Palamareva, M. and Kozekov, I.
Chromatographic behaviour of diastereoisomers. XIII. Adsorptivity of esters of *Z*- and *E*-2,3-diphenylpropenoic acids and similar compounds on silica in terms of the Snyder theory 758(1997)135
Påhlsson, P., see Ohlson, S. 758(1997)199
Pena, A.L.S. and Silveira, M.I.N.
Determination of chlorophenoxy herbicides in waters by capillary gas chromatography with ion trap detection 758(1997)324
Perrin, S.R., see Welch, C.J. 758(1997)93
Pişkin, E., see Denizli, A. 758(1997)217
Polinina, E., see Fairman, B. 758(1997)85
Popl, M., see Sýkora, D. 758(1997)37
Richard, J.-M., see Oubrahim, H. 758(1997)145
Richmond, R.
Database of structures and their gas chromatography retention indices, tagged with individual search windows 758(1997)319
Ritter, C., see Gilliard, J.A. 758(1997)1
Runnels, P.L., see Weldon, M.K. 758(1997)293
Saito, K., see Kiyohara, S. 758(1997)209
Salih, B., see Denizli, A. 758(1997)217
Seigle-Murandi, F., see Oubrahim, H. 758(1997)145
Silveira, M.I.N., see Pena, A.L.S. 758(1997)324
Simonetti, M.S., see Damiani, P. 758(1997)109
Smith, R.M., see Zimina, T.M. 758(1997)191
Sugita, K., see Kiyohara, S. 758(1997)209
Sugo, T., see Kiyohara, S. 758(1997)209
Susic, M. and Alongi, D.M.
Determination of terrestrial markers in marine environments by gas chromatography–mass-selective detection compared to high-performance liquid chromatography–fluorescence detection 758(1997)243
Sýkora, D., Tesařová, E. and Popl, M.
Interactions of basic compounds in reversed-phase high-performance liquid chromatography. Influence of sorbent character, mobile phase composition, and pH on retention of basic compounds 758(1997)37
Szczerba, T., see Welch, C.J. 758(1997)93
Tesařová, E., see Sýkora, D. 758(1997)37
Toida, Y., see Kiyohara, S. 758(1997)209
Tomás-Barberán, F.A., see Montijano, H. 758(1997)163
Torre, M., see García, M.C. 758(1997)75
Trécourt, F., see Oubrahim, H. 758(1997)145
Volgger, D., Zemann, A.J., Bonn, G.K. and Antal, Jr, M.J.
High-speed separation of carboxylic acids by co-electroosmotic capillary electrophoresis with direct and indirect UV detection 758(1997)263
Wan, H. and Blomberg, L.G.
Enantiomeric and diastereomeric separation of di- and tripeptides by capillary electrophoresis 758(1997)303
Welch, C.J., Szczerba, T. and Perrin, S.R.
Some recent high-performance liquid chromatography separations of the enantiomers of pharmaceuticals and other compounds using the Whelk-O 1 chiral stationary phase 758(1997)93
Weldon, M.K., Arrington, C.M., Runnels, P.L. and Wheeler, J.F.
Selectivity enhancement for free zone capillary electrophoresis using conventional ion-pairing agents as complexing additives 758(1997)293
Wheals, B., see Fairman, B. 758(1997)85
Wheeler, J.F., see Weldon, M.K. 758(1997)293
Yun, T., Bensetiti, Z., Zhong, G. and Guiochon, G.
Effect of column efficiency on the internal concentration profiles and the performance of a simulated moving-bed unit in the case of a linear isotherm 758(1997)175
Zemann, A.J., see Volgger, D. 758(1997)263
Zhong, G., see Yun, T. 758(1997)175
Zimina, T.M., Smith, R.M. and Myers, P.
Comparison of ODS-modified silica gels as stationary phases for electrochromatography in packed capillaries 758(1997)191